

Horticulture

extension.usu.edu

March 2018

Horticulture/Fruit/2018-02pr

Selecting Blackberry Cultivars for Utah

Brent Black, Thor Lindstrom, Britney Hunter, Shawn Olsen, Sheriden Hansen, and Tiffany Maughan

Introduction

Historically, Utah has not been a significant blackberry producer. This is likely due to harsh winters and frequent late spring frosts, which result in significant crop loss. Blackberry canes often suffer from some cold injury in Northern Utah, which typically results in loss of floricanes buds, or dieback of the entire cane (Figure 1), although Utah winter conditions rarely cause permanent damage to the plant.

Despite their cold sensitivity, the delicate fruit have a short shelf life that makes shipping the fruit from distant markets difficult, presenting opportunities for local sales. With careful cultivar selection and good management, blackberries can be successfully grown in Utah and can be a profitable fresh-market crop.

For 6 years, 21 blackberry cultivars were compared at the USU Kaysville Research Farm (Kaysville, Utah) in order to identify cultivars that might be suited to local production. Characteristics evaluated were winter survival, yield, and fruit quality suitable for direct-market sales.

Fruiting Type

There are two fruiting types of blackberries: floricanes and primocane. Floricane-fruiting blackberries bear fruit only on second-year canes. After fruiting, these canes are pruned out leaving the first-year canes to produce the next year. Primocane-fruiting blackberries produce fruit on first-year canes, an advantage in that the canes do not have to survive a winter. These types are pruned to the ground each year.

Growth Habit

Blackberries are classified by growth habit into three categories: trailing, semi-erect, and erect. Trailing types produce long thin canes that will grow along the ground unless supported by an appropriate trellis. Semi-erect

blackberries have primocanes that initially grow erect, but then branch and the side branches droop down to the ground. Erect blackberries are mostly self-supporting plants but still benefit from trellising.

For more detailed information on fruiting types and growth habits of blackberries reference our [Blackberry Management](#) fact sheet.



Figure 1. Winter injured floricanes with new primocanes emerging.

Kaysville Cultivar Trial

Blackberry plants from 21 different cultivars and advanced selections were established in 2006. Plants were spaced 5 feet within the row and rows were spaced 10 feet apart. Irrigation was provided using both drip and overhead systems. Half of the replications were placed on a vertical trellis; the other half were placed on a rotating cross-arm trellis. Primocanes were selected and tied to the trellis periodically throughout the growing season. The rotating cross-arm was set near the ground in December each year. All floricanes were pruned the winter following fruiting.

In the 2008-2012 growing seasons, plots were evaluated for winter injury, yield, fruit size, and production season. Once each week, a subsample of fruit was taken and weighed to determine average fruit size. Ripe fruit was harvested three times per week from mid-July until the first fall freeze. Consumer taste tests were carried out in 2011 and 2012.

Results

WINTER INJURY

Based on a visual assessment of the percent of surviving buds, winter injury was compared each spring. One particularly hard winter was noted in the study, 2010-2011. Temperatures dropped from 50°F to -0.4°F in 48 hours on November 24-25, without any prior acclimating temperatures. See Table 1 for winter survival of each cultivar over 6 years. 'Illini Hardy' (average 93% survival), 'Apache' (88% survival), and 'Chester' (83% survival), consistently had the highest rate of winter bud survival. By contrast, 'Marion' (19% survival), and 'Kiowa' (16% survival), had the poorest rates of bud survival. There was no significant difference in bud survival rates between the vertical trellis and the swing trellis reps. Note that winter injury is not reported for 'Prime Jim' and 'Prime Jan' (Primocane-fruited) since canes are removed to the ground each fall.

YIELD

The cultivars with the largest fruit size included 'Kiowa', 'Triple Crown', and 'Apache'. The cultivars 'Navaho' and 'Chester' had the smallest fruit size. Yields were closely correlated with winter survival. 'Triple Crown' consistently had the highest yields, followed by 'Doyle's' and 'Hull'. The trailing types ORUS 1939-4, 'Metolius', and 'Marion' consistently had the lowest yields.

Another way to evaluate yield is to look at how reliably the cultivar will produce a crop each year. Since yield varies considerably between years, a reliability index (75%) was calculated. This represents the minimum yields that could be expected in 75% of the years. Yield reliability is shown as a black line on the bar graphs (Figure 2). In general, only the semi-erect types had yield reliability at acceptable levels for Utah cultivation (ranging from 0.3 to 1.0 pounds per plant depending on cultivar). Trailing types had yield reliability indexes less than zero, indicating that total crop failure would result more than 25% of the time.

CONSUMER PREFERENCE

In order to gauge consumer preference, some cultivars were evaluated in taste tests at the USU Botanical Center farmers market in Kaysville in 2011 and 2012. Cultivars

Table 1. Average winter survival over 6 years: 2007-2012, and average fruit size (gram/fruit) over 5 years: 2008-2012.

	% Survival	grams/fruit
Illini Hardy	92.2	3.1
Apache	87.7	5.4
Chester	84.7	2.9
Navaho	78.5	2.9
Triple Crown	76.9	5.7
Arapaho	75.0	4.3
Hull	66.2	4.2
Ouachita	56.7	4.8
Doyle's	58.3	3.1
Newberry	57.5	4.3
Loch Ness	61.3	3.3
Siskyou	52.5	4.0
Black Diamond	32.0	3.5
Obsidian	25.1	4.1
ORUS 1793-1	27.0	4.6
ORUS 1939-4	26.5	3.3
Metolius	25.5	3.9
Marion	14.0	3.2
Kiowa	17.3	6.7
Prime Jim	--	4.3
Prime Jan	--	4.4

were placed in clamshells with a numbered piggybank behind each cultivar. Market patrons were given 10 pennies and instructed to put one or more pennies in the piggybanks adjacent to their favorite cultivars. Due to differences in ripening, not all cultivars were available for all tastings. Six cultivars were included in all three tastings, and a seventh cultivar was present at two of the three tastings. These cultivars are compared in Table 2. The top four cultivars in tastings over 2 years based on relative order of preference were: 'Apache', 'Arapaho', 'Triple Crown', and 'Chester'. 'Arapaho' was not available for tasting on 8-Sep-2011.

Table 2. Taste preferences ranking based on overall votes for each tasting date. Rank was calculated based on highest to lowest total "votes" (number of pennies) for taste preference with rank 1 receiving the most votes.

	8/18/11	9/8/11	8/9/12	Avg.
Apache	3	1	1	1.7
Arapaho	2		2	2.0
Triple Cr.	1	3	3	2.3
Chester	5	2	5	4.0
Doyle's	7	5	4	5.3
Loch Ness	6	4	8	6.0
Illini Hardy	8	6	7	7.0




Figure 2. Average yield with yield reliability index (75%, represented by black line) of 21 cultivars at the Kaysville Research Farm. Values are average of five seasons (2008-2012). Yield reliability index was below 0 for all trailing type cultivars so no line appears in the graph.

Cultivar Descriptions

SEMI-ERECT TYPES:

‘Triple Crown’ - (Thornless) This is a long-time favorite among Utah blackberry growers for its productivity, vigor and flavor. Often considered the standard for comparison among blackberries, it produces large berries mid-season and has heavy yields. It was susceptible to moderate winter damage but was a favorite among taste test consumers. Developed by the USDA Agricultural Research Service.

‘Doyle’s Thornless’ - (Thornless) Heavy producer, but fruit size was among the smallest. Produces mid-season with moderate winter survival. Not a favorite among consumers.

‘Hull’ - (Thornless) Good winter survival rate, produces medium sized fruit mid-to-late season. No taste test data on this variety. It was introduced in 1981 from Illinois Agricultural Experiment station. Fruit are oblong and color well in the heat.

‘Chester’ - (Thornless) Exceptional winter survival. Can fruit quite late in the season and will produce high yields with mild fall weather, but may freeze out before full yield potential is reached. Fruit size was the smallest of the cultivars tested, and the fruit was not favored by consumers. Developed by the USDA Agricultural Research Service.

‘Loch Ness’ - (Thornless) This moderate yielding plant produces medium sized fruits mid-season. Winter survival is fair. This was not a favorite among taste testers.

ERECT TYPES:

All erect types, except Illini Hardy were released from the University of Arkansas breeding program.

‘Illini Hardy’ - (Very thorny) Developed by the University of Illinois for northern growers. Best rate of winter survival of all the cultivars in the trial. This plant produces small fruit late in the season with heavy yields. Good for areas with late frosts and/or unpredictable springs. Least favorite among taste tested cultivars.

‘Arapaho’ - (Thornless) produces medium to large fruit mid to late season (ripens before ‘Apache’ and ‘Navaho’). It was among the more winter hardy, and the highest yielding of the erect types. This was a taste test favorite among consumers.

‘Ouachita’ - (Thornless) This cultivar produces large fruit late in the season, with moderate yields and good postharvest storability. ‘Ouachita’ scored low in consumer taste tests.

‘Apache’ - (Thornless) A favorite among consumers. Produces moderate yields of large fruits in mid-season. Most winter hardy of the erect types.

‘Navaho’ - (Thornless) Produces below-average yields, with fruit size among the smallest tested. It had very good winter survival. Fruit was not included in taste tests.

‘Kiowa’ - (Thorny) Least winter hardy variety of all cultivars tested with an average of 16% cane survival. It was low yielding but had extremely large fruit. No taste test data available.

PRIMOCANE-FRUITING TYPES (erect):

‘Prime Jim’ - (Thorny) Developed at the University of Arkansas. Plants bear fruit on first year canes. This cultivar produces larger fruit than ‘Prime Jan’ slightly earlier in the season. This cultivar also produced heavier yields than ‘Prime Jan’. No winter damage assessment was taken. No taste test data are available.

‘Prime Jan’ - (Thorny) Plants bear fruit on first year canes, eliminating problems associated with winter cane survival. Plants produced moderate harvests late season. Fruits were moderately sized. No taste test data are available. Released from the University of Arkansas.

TRAILING TYPES:

All trailing types were developed in Corvallis, Oregon, by the USDA – Agricultural Research Service.

‘Newberry’ - (Thorny) Moderate winter survival. This plant produces medium-to-large fruit early in the season. It had the best winter survival and the highest yields

among trailing types, and had moderate scores in taste tests.

‘Siskyou’ - (Thorny) A favorite among consumers participating in the taste tests, it produced large fruit, and was the earliest bearing cultivar in the planting. Yields were inconsistent from year to year due to winter damage.

‘Obsidian’ - (Thorny) Produces medium-sized fruit mid-season. It has poor winter survival and very inconsistent yields. No taste test data available.

‘Marion’ - (Thorny) This cultivar is widely used for commercial production and is typically used in processing. It is not well-suited to the climate in northern Utah.

‘Black Diamond’ - (Thornless.), **‘Metolius’** - (Thorny.), **ORUS 1793** - (Thorny.), and **ORUS 1939** - (Thorny.) were not sufficiently winter hardy for northern Utah.

This project is funded in part by USDA-Risk Management Agency under a cooperative agreement. The information reflects the views of the author(s) and not USDA-RMA.



Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran’s status. USU’s policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran’s status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Ken White, Vice President for Extension and Agriculture, Utah State University. (3-2018)